

Outpatient therapy now available for those with COVID-19. Supplies may be limited.

Eligibility guidelines ensure those with the highest risks have priority access to treatments.

Those at highest risk	Treatment window after symptoms appear:	Paxlovid PO	Sotrovimab IV	Remdesivir IV	Molnupiravir PO
		5 days	10 days	7 days	5 days
		✓	✓	✓	✓ <i>If other therapies not available or appropriate</i>
		✓	✓	✓	✓ <i>If other therapies not available or appropriate</i>
		✓	✓	✓	✓ <i>If other therapies not available or appropriate</i>
		✓	Not currently eligible	Not currently eligible	✓ <i>If other therapies not available or appropriate</i>
		Not currently eligible	Not currently eligible	Not currently eligible	✓
		Not currently eligible	Not currently eligible	Not currently eligible	✓

*Up to date means a person has received all recommended COVID-19 vaccines, including booster dose(s) when eligible (bit.ly/CDCStayUptoDate).

**MI priority risk factors include:
Obesity (Body Mass Index >35), chronic respiratory disease, pregnancy (note: in pregnancy, molnupiravir should not be used and Paxlovid and remdesivir should be used with caution when sotrovimab is unavailable), chronic kidney disease (special considerations with Paxlovid), cardiovascular disease, and diabetes.



Talk to your health care provider or visit Michigan.gov/COVIDtherapy to learn more.